



We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.

To use the Text Relay service, prefix all numbers with 18802.

اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید۔

ने उदाहृत उदाहरणों में लैंग्वेज व अंतरांतरण से इस संख्या से बात करें: 01932 723553

اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براہ کرم اس فون نمبر 01932 723553 پر رابطہ کریں

Se precisa de uma tradução por favor contacte: 01932 723553

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553

यदि आपको अनुवाद की ज़रूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553

Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

Ashford Hospital
London Road
Ashford, Middlesex
TW15 3AA
Tel: **01784 884488**

St. Peter's Hospital
Guildford Road
Chertsey, Surrey
KT16 0PZ.
Tel: **01932 872000**

Website: www.ashfordstpeters.nhs.uk

Ankle Injury

Paediatric A&E Department

Ankle Injury

Your child has been treated for an injury to the ankle. When you are at home the following will help:

1. Elevate the foot to above the level of the heart when resting by using pillows or cushions; this will help to reduce the swelling.
2. Use an ice pack to ease pain, swelling and bruising. You can use frozen peas or crushed ice, but whichever you choose ensure it is wrapped in a damp towel first.

Never apply an ice pack directly to the skin.

An ice pack should be placed around the elevated ankle for no more than 10 minutes and this can be repeated 4 - 6 times a day.

The ankle should be completely rested for the first 24 hours after the injury. When your child starts to walk on the injured ankle ensure that they take even steps placing the heel down first and then the toe. Only put as much weight onto the foot as pain allows and use crutches or a stick if they have been advised to do so.

EXERCISES

Exercises should be done for ten minutes once an hour, especially after ice treatment to avoid the ankle stiffening up. Repeat the following ten times each.

1. Point the foot upwards and then point it down again, wiggle the toes.
2. Gently rotate the ankle in a circle keeping the knee straight.
3. Place feet side by side with ankles together. Then turn the soles of both feet first towards each other and then away from each other.

Give your child a medicine such as Calpol, Ibuprofen or Paracetamol to relieve any pain. Please follow the dosage and frequency instructions on the bottle / packet carefully.

If the pain increases or the swelling gets worse contact your GP for advice.

If you have been seen in the Accident and Emergency Department in the evening or a weekend and you have been given a Physiotherapy card for your child, please attend the Physiotherapy Department between 09.00 - 10.00 a.m. on the next working day.

FURTHER INFORMATION

Paediatric Accident and Emergency Department
St. Peter's Hospital
Guildford Road
Chertsey
Surrey KT16 0PZ
Telephone **01932 872000**

For reassurance and advice please contact NHS Direct on: **0845 4647**

The following websites also gives useful advice: www.patient.co.uk and www.kidshealth.org

Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty. If they cannot resolve your concern, please contact our Patient Advice and Liaison Service (PALS) on 01932 723553 or email pals@asph.nhs.uk. If you still remain concerned please contact our Complaints Manager on 01932 722612 or email complaints@asph.nhs.uk.

Author: Dr Erin Dawson

Department: Paediatric A&E

Version: 2

Published: July 2008

Review: July 2011